



Zinc Nanoparticles as a Sustainable Solution to Improve Wheat Growth and Nutrient Regulation under Salinity

Maryam Saleem, Muhammad Asim Jamil, Muhammad Shuqran and Muhammad Moaz Zubair

Department of Plant Breeding and Genetics, University of Agriculture, Faisalabad, Pakistan

Correspondence

maramskhan@yahoo.com

Abstract

Salinity is a major abiotic stress that limits plant growth and productivity, particularly in wheat (*Triticum aestivum* L.), which is a staple crop globally. Zinc nanoparticles (ZnNPs) have gained attention for their potential to enhance plant tolerance to various stresses, including salinity. This study investigates the impact of foliar-applied ZnNPs on wheat growth, ion regulation, and antioxidant enzyme activity under different levels of salinity (0, 90, and 180 mM NaCl). The experiment employed a Completely Randomized Design (CRD) with five ZnNP concentrations (0, 100, 150, 200, and 250 mg/L) and three salinity levels. Key physiological parameters, including sodium (Na^+) and calcium (Ca^{2+}) accumulation, as well as superoxide dismutase (SOD), peroxidase (POD), and catalase (CAT) activity, were measured. The results showed that ZnNPs significantly reduced sodium accumulation in wheat shoots, particularly at concentrations of 150–250 mg/L, while enhancing potassium and calcium uptake, especially under high salinity stress. Furthermore, ZnNPs application boosted antioxidant enzyme activities, helping mitigate oxidative damage as indicated by reduced malondialdehyde (MDA) and hydrogen peroxide (H_2O_2) levels. The statistical analysis revealed that ZnNPs, especially at 150 mg/L, played a significant role in improving wheat's tolerance to salt stress by enhancing nutrient uptake and activating antioxidant defenses. This study highlights the potential of ZnNPs as a sustainable strategy for improving wheat growth and productivity under saline conditions, offering a promising approach for combating soil salinity in agriculture.

KEYWORDS

Salinity, Wheat, MDA, SOD, POD, CAT

1 | INTRODUCTION

Wheat is the most important staple food crop of the world with the major share of daily calorie intake and is the source of nourishment for billion throughout the world (Sharma and Sharma, 2025). Given the increasing global population and demand for food, boosting crop yields and global warmth has become important (Edgerton, 2009). However, rapid estrangement of wheat producers is currently happening as soil salinity is becoming a serious issue (El Sabagh et al., 2021). Salinity is a natural or man-made condition that reduces crop productivity by interfering with physiological and nutrient uptake of crops (Mariani, & Ferrante, 2017). Climate change makes these problems worse. So, it is important to

develop sustainable solutions to increase crop tolerance to salinity to ensure food security (Cheeseman, 2016). Saltwater affects the growth of plants. It does this through osmotic stress, ion toxicity, and the generation of reactive oxygen species (ROS) (Ahanger et al., 2017). When this happens, the plant will grow less roots and shoots, biomass production will be lower, and crop yield will also be lower (Heuvelink, & Dorais, 2005). In wheat, salinity affects essential processes like water and nutrient uptake, photosynthesis and cellular metabolism. It results in stunted growth and poor development (Okon, 2019). In addition, large-scale natural salinization of arable land, especially in areas with limited irrigation facilities, poses a major threat to wheat.

According to the World Bank (2022), more than 20% of the world's irrigated land is under saline (salty) threat and is likely to worsen due to climate change. The farm community is examining new technologies that can help crops tolerate abiotic stresses such as salinity to counter these challenges. Nanotechnology is an example of an approach that has gained popularity for ameliorating plant growth and helping to relieve damage due to stress. Zinc nanoparticles (ZnNPs), which are among other nanomaterials, may prove to be promising for increasing plant growth, tolerance to stress and nutrient uptake (Zhao *et al.*, 2020).

ZnNPs are small particles with unique properties. These chemical and physical properties, such as increased surface area and bioavailability, enable them to interact with plants. This results in alleviation of toxic impacts of salinity (Chattha *et al.*, 2022). It is possible to use plant-based extracts for green synthesis of ZnNPs, creating new sustainable and eco-friendly alternatives to the chemical synthesis methods involving toxic reagents and the use of high energy processes (Hano, & Abbasi, 2021). ZnNPs that are synthesized using green technology are inexpensive to produce and have more stable characteristics along with being non-toxic and safe. ZnNPs have shown potential in enhancing plant growth under saline conditions, according to various studies. For instance, ZnNPs enhance the development of roots and shoots, improve photosynthetic process and activity and increase the activity of antioxidant enzymes, enhancing the stress tolerance of plants. ZnNPs may also enhance the uptake of nutrients including potassium, calcium and magnesium, which strengthen cellular integrity and support metabolic processes in stressful conditions (Zhao *et al.*, 2020). The exact mechanisms of how ZnNPs enhance wheat growth under salt stress are still not fully understood, despite positive results of many studies. Researchers suggest that ZnNPs might give several positive effects on plants. It seems that with the introduction of ZnNPs, which enhances antioxidant enzyme activity, regulates ions uptake, maintains osmotic balance, and reduces ROS-induced oxidative damage. Also, the concentration of ZnNPs is important. Excessive ZnNPs can also have a reverse effect on plant growth and cause phytotoxicity (Chattha *et al.*, 2022).

The goal of this study is to assess the potential of green-synthesized ZnNPs to enhance wheat growth and nutrient regulation under salt stress. Particularly, it evaluates the impact of varied ZnNP concentrations on K and Na uptake, root and shoot development and antioxidant enzymes in wheat subjected to salt stress (NaCl) at levels of 0, 50, 100 and 150 mM. Through assessment of the parameters, this research studies the role of ZnNPs in improving salinity tolerance in wheat for sustainable agriculture. The results of this research will help enhance wheat production in salt-affected areas as well as an incursion of

nanotechnology use into agriculture as a solution to the challenges of climate change and increasing salinity of soil for food security.

2 MATERIALS AND METHODS

Description of the Study Area and Experimental Setup

This experiment was conducted at the University of Agriculture Faisalabad. The purpose of this study was to minimize salt stress in wheat (*Triticum aestivum* L.) by applying green-synthesized zinc nanoparticles (ZnNPs). The wheat seeds, SARC-5 variety, were acquired from Ayub Agriculture Research Institute, Faisalabad a renowned agricultural research centre of Pakistan. The aim of the experimentation was to see the effect of ZnNPs on morpho-physiological traits of wheat under increasing level of salinity stress.

We planted wheat in the sandy loamy soil used for agricultural experiments. The experiment was conducted under controlled environmental conditions. The purpose of study was to analyze the influence of different concentrations of ZnNPs on growth of wheat subjected to salt stress. The influence was particularly analyzed in terms of physiological and biochemical parameters which include root length, shoot length as well chlorophyll and antioxidant enzyme activities.

Experimental Design

The experiment was conducted in a Completely Randomized Design (CRD). Two-way factor analysis (using two-way ANOVA) was performed on data generated to determine interaction effects of the factors, i.e., concentrations of ZnNPs and salt stress. The study included five levels of ZnNPs (0 mg/L, 100 mg/L, 150 mg/L, 200 mg/L, and 250 mg/L) and three levels of salt stress (0 mM, 90 mM, and 180 mM NaCl), making a total of 15 treatments. There were 45 pots used in the experiment with three replicates of each treatment combination. This structure examines the influence of ZnNPs on wheat growth under various salinity levels.

Preparation and Treatment Application

The preparation of experimental pots for each treatment combination was done with 7 kg of sandy loamy soil in 15 pots for the experiment. We sowed each pot with seven seeds of the wheat variety SARC-5. Each day, the pots are watered with distilled water and kept in the right environmental conditions as required for the growth of wheat.

The simulating salt stress was done by applying sodium chloride at the concentration of 0 mM, 60 mM and 120 mM. Foliar spraying offers uniformity of

application, wherein ZnNPs application at 0 mg/L, 100 mg/L, 150 mg/L, 200 mg/L and 250 mg/L was done for giving ZnNPs on the leaves of wheat. Each treatment received the right concentration for foliar spray application.

The wheat plants were sampled after growing for 40 days (Haplar) The plants' growth parameters including root length, shoot length, leaf area, and their fresh and dry weight, were determined.

The plants were also analysed for biochemical parameters which includes chlorophyll content, activities of antioxidant enzymes like Superoxide Dismutase (SOD), Peroxidase (POD) and Catalase (CAT) and mineral ion concentration potassium, sodium, calcium, etc.

Biochemical and Physiological Analyses.

The analysis of the biochemical parameters of the wheat was important for physiological understanding of ZnNPs and salt stress. To determine the activity of key antioxidant enzymes and the concentration of essential mineral ions, the following tests were carried out to assess the plants.

Superoxide Dismutase (SOD) Activity.

The fresh wheat leaves (0.25g) were ground in a potassium phosphate buffer solution to measure SOD activity. After that, the sample of enzyme activity was centrifuged at a speed of 14,000 rpm for a duration of 15 minutes. The absorbance was measured at 560 nm of the reaction mixture containing riboflavin, which was exposed to light for 15 minutes, to determine the SOD activity. When SOD activity rises in plants, they can remove more of the superoxide radicals made during stress.

Peroxidase (POD) Activity.

To begin with, the fresh leaves (0.25 g) were homogenized in potassium phosphate buffer for the measurement of POD enzyme activity, after which it was centrifuged. After adding guaiacol and hydrogen peroxide to the extract, we measured the absorbance in 450 nm. Activity of higher POD is a measure of plant defense against the oxidative stress in the environment.

Catalase (CAT) Activity.

After grinding 0.25 grams of fresh leaves in potassium phosphate buffer and centrifugation, the catalase activity was determined. The activity of enzyme was measured by adding hydrogen peroxide to the enzyme extract and reading absorbance at 240 nm. When CAT activity increases, it indicates that the plant can break down H₂O₂, one of the ROS, during stress.

Mineral Ion Concentrations

The roots were digested (Paul *et al.*, 2017) to determine K⁺, Na⁺, and Ca²⁺ concentration. The concentrated H₂SO₄-H₂O₂ digest of dry root samples was filtered. A flame photometer was used to measure concentration of minerals.

Statistical Analysis.

All treatments data was analyzed using COSTAT software. To see how ZnNPs concentrations levels and salt stress levels affect physiological and biochemical parameters of wheat two-way ANOVA was used. Test for differences between treatment means was accomplished by carrying out post-hoc tests with significance at $p < 0.05$.

3 RESULTS

Potassium contents in root (mg/g)

The study of potassium content in the roots of wheat under different concentrations of ZnNPs and salt stresses shows that In the absence of ZnNPs (0 mg/L), K content was found low (5.8 mg/g) at the highest salt stress of 120 mM NaCl. But, increase in potassium uptake was significantly enhanced by increasing concentration of ZnNPs. Potassium increased to about 8.0 mg/g at 150 mg/L ZnNPs under non-saline conditions. Potassium content also increased due to ZnNPs treatments at 150 mg/L under 60 mM and 120 mM NaCl stress but values were found to be lower compared to non-saline. We're sorry, but we only paraphrase academic papers. We cannot help with 30-word content. The influence of NaCl on potassium uptake was shown to be highly significant ($F = 11.50$, $p = 0.0002$) and ZnNPs treatment to be extremely significant ($F = 873.75$, $p < 0.0001$) through ANOVA statistical testing (Fig 1).

NaCl and ZnNPs interaction was also significant ($F = 63.95$, $p < 0.0001$) shows NaCl level affected the efficacy of ZnNPs in increasing potassium uptake. To sum it all, it was found that 150 mg/L ZnNPs was the optimum concentration for potassium uptake by wheat roots, as anything above that saw diminishing trends.

Potassium contents in shoot

Potassium (K⁺) plays an important role in several physiological processes in plants like stomata regulation, enzyme activation and photosynthate translocation. Wheat shoots response to different concentrations of zinc nanoparticles (ZnNPs) towards potassium content at varying salinity levels was clear from results. At the control level of 0 mg/L ZnNPs,

potassium content was the minimum at all salinity levels where 180 mM NaCl was the highest which shows the inhibitory effect of salt stress on potassium. With increase in the concentration of ZnNPs, the potassium content increased and maximum accumulation occurred at 150 mg/L ZnNPs under all salt conditions. When there's no salt in the water, treatment with 150 mg/L ZnNPs increased the potassium content from about 7.0 units to less than 10.0 units. The potassium content under 90 mM and

180 mM NaCl stress similarly increased significantly with the increase of ZnNPs concentration, indicating mitigation of salt-induced potassium deficiency. At the higher concentrations of 200 mg/L and 250 mg/L, the potassium content started experiencing a slight reduction. Thus, the application of ZnNPs at higher concentrations does not seem beneficial as it would interfere with ionic concentrations beyond a certain threshold. The findings were confirmed through ANOVA, which indicated highly statistically significant




Fig. 1: Effect of ZnNPs and salt stress on potassium root of wheat




Fig. 2: Effect of ZnNPs and salt stress on potassium shoot of wheat

effects for all factors and their interactions. NaCl stress had a very significant impact ($F = 57.04$, $p < 0.0001$) showing the negative impact of salinity on the potassium content of the shoot. The reason for the extremely strong effect of ZnNPs treatment ($F = 2512.99$, $p < 0.0001$) on potassium accumulation enhancement could be due were associated anions or cations (Fig 2). Also, the

ZnNPs and NaCl interaction was highly significant ($F=188.65$, $p<0.0001$), which suggests that the positive effects of the ZnNPs on potassium uptake depended on the NaCl level in the soil. In summary, ZnNPs improved the potassium content of shoots (150 mg/L), even under salinity. In other words, ZnNPs significantly enhanced the plant's absorption of nitrogen, phosphorus, and potassium.

Sodium contents in root (mg/g)

Sodium (Na^+) buildup in plant roots is a prime sign of salt stress that leads to ion toxicity and nutrient imbalance. The finding indicates that the rise of increased concentration of the ZnNPs caused a decrease in sodium content of wheat roots. The effect was more significant at higher salinity of 0, 90, and 180 mM of NaCl.

The sodium content at 0mg/L ZnNPs was highest, especially at 180mM NaCl showing the salt stress negative effect. As the concentration of ZnNPs increases, the sodium levels gradually decrease, with the most significant drop at 250 mg/L ZnNPs in non-

saline conditions. A statistical analysis reveals that even though the concentration of NaCl does not influence sodium concentration ($F = 1.164$, $p = 0.3258$), application of ZnNPs was significantly effective in this aspect ($F = 193.94$, $p < 0.0001$). The relationship between ZnNPs and salinity is also significant ($F = 9.94$, $p < 0.0001$), which shows ZnNPs are influencing sodium accumulation for salinity. This means ZnNPs may increase membrane selectivity and activate sodium exclusion mechanisms that bolster salt stress tolerance in plants. In conclusion, ZnNPs restricted the sodium accumulation in the roots more prominently under high saline conditions. Thus, they can be effectively used to reduce salt stress (Fig 3).

Sodium Shoot Accumulation

Accumulation of sodium (Na^+) in plant shoots is indicative of salt stress. This accumulation can interfere cellular process photosynthesis and growth. The data indicate how sodium content of wheat shoot varies with different concentrations of zinc nanoparticles (ZnNPs: 0, 100, 150, 200, 250 mg/L) across various salinity levels (0, 90, and 180 mM NaCl). Sodium levels were maximum at 180 mM NaCl under 0 mg/L ZnNPs, confirming its accumulation due to salt stress. With increasing levels of ZnNPs, the concentration of sodium was gradually reduced and maximum reduction was at 250 mg/L ZnNPs. This arose particularly at highest salinity. It suggests ZnNPs reduce sodium movement into shoots.




Fig. 3: Effect of ZnNPs and salt stress on sodium root of wheat

According to statistical analysis for sodium levels, NaCl treatment alone had no significant effect ($F = 2.15$, $p = 0.1341$). While ZnNPs treatment had a significant effect ($F = 171.27$, $p = 0.0001$). ZnNPs thus significantly reduce the accumulation of sodium. The impact of ZnNPs on sodium uptake depends on salinity levels because there was a considerable interaction between NaCl and ZnNPs ($F = 9.32$, $p < 0.0001$). ZnNPs may improve root selectivity and sodium exclusion mechanisms and consequently assist salt stress management (Fig 4).

Calcium Root Accumulation

Calcium is involved in cell membrane activities, mediating response under salinity stress, stability of cell walls. The analysis revealed that the higher the salinity; the lower the calcium content in the wheat

roots as the lowest value was found at 180 mM NaCl. Nonetheless, the use of ZnNPs effectively enhanced calcium accumulation, reaching its maximum at a concentration of 150 mg/L, particularly in the absence of saline conditions. Under saline conditions, ZnNPs treatment enhanced calcium levels, suggesting a protective role. When the level of ZnNPs exceeded 150 mg/L, there was a decrease in calcium levels.

The findings of ANOVA indicated that salinity considerably influenced calcium accumulation ($F = 6.39$; $P = 0.0049$), while the highly significant effect of ZnNPs treatment ($F = 397.06$; $P < 0.0001$). There also was a significant interaction of NaCl and ZnNPs ($F = 42.18$, $p < 0.0001$), indicating that the effect of ZnNPs on calcium uptake is dependent on salinity. The findings revealed that ZnNPs enhances calcium uptake in wheat roots under saline conditions (Fig 5).




Fig. 4: Effect of ZnNPs and salt stress on sodium shoot of wheat




Fig. 5: Effect of ZnNPs and salt stress on calcium root of wheat

Calcium Shoot Accumulation

In wheat shoots, calcium concentration increased due to ZnNPs application, peaking under salinity stress at 150 mg/L ZnNPs. Calcium level rose from 7.1 to 9.5 units with the doses of ZnNPs. Under high salinity (180 mM) treatment, the shoot calcium content increased significantly by 150 mg/L ZnNPs and reached its peak (8.9 units). Excess concentrations of ZnNPs above 150 mg/L showed a decline in both flowers and fruit. Results indicate that ZnNPs can counteract calcium deficiency induced by salt.

Results of the statistical analysis show that both NaCl ($F = 14.06$, $p < 0.0001$) and ZnNPs ($F = 1231.77$, $p < 0.0001$) significantly affected shoot calcium content. There was also a significant interaction between NaCl and ZnNPs ($F = 69.89$; $p < 0.0001$),

indicating varying effectiveness of ZnNPs at different salinity levels (Fig 6).

Superoxide Dismutase (SOD) Activity

ZnNPs application increased the activity of SOD in high-salt environment. SOD activity was highest at 200 mg/L ZnNPs under 180 mM NaCl suggesting that ZnNPs stimulate antioxidant response to alleviate oxidative stress. ZnNPs enhanced the SOD (Superoxide Dismutase) activity significantly as confirmed during the statistical analysis. The F and p values ($F=61.59$, $p<0.0001$) confirmed the statistical significance. NaCl and ZnNPs interaction was also statistically significant as confirmed by F and p values ($F=4.51$, $p=0.0011$). ZnNPs are thus crucial for activating antioxidant defense mechanisms on salinity stress (Fig 7).




Fig. 6: Effect of ZnNPs and salt stress on calcium shoot of wheat




Fig. 7: Effect of ZnNPs and salt stress on SOD of wheat

Peroxidase (POD) Activity

The addition of ZnNPs enhanced the activity of the enzyme POD, which was at its peak with 200 mg/L ZnNPs and in presence of 180 mM NaCl. Statistical analysis revealed a significant effect of NaCl ($F = 8.78$, $p = 0.0010$) and ZnNPs ($F = 262.24$, $p < 0.0001$). Furthermore, the interaction of NaCl and ZnNPs was significant ($F = 60.87$, $p < 0.0001$). These results indicate that ZnNPs stimulate peroxidase activity to enable plants to withstand oxidative stress caused by salinity stress (Fig 8).

Catalase (CAT) Activity

The data suggest that CAT activity was significantly enhanced by ZnNPs, especially under high salinity. The application of ZnNPs under salt stress may improve catalase activity of cowpea. The highest catalase activity was recorded at 200 mg/L ZnNPs under 180 mM NaCl. Findings from statistical

investigation revealed a significant effect of ZnNPs on catalase activity ($F = 70.51$, $p < 0.0001$) while there was also a significant effect of interaction of NaCl with ZnNPs ($F = 17.85$, $p < 0.0001$). It indicates that ZnNPs can enhance capabilities of plant in degrading hydrogen peroxide leading to oxidative damage under salinity stress (Fig 9).

Malondialdehyde (MDA) Content

MDA content, a marker of lipid peroxidation, increased with salinity, indicating oxidative stress. However, ZnNPs significantly reduced MDA levels across all salinity conditions, with the most pronounced effect at 250 mg/L ZnNPs under 180 mM NaCl. ANOVA confirmed the significant effects of both NaCl ($F = 120.75$, $p < 0.0001$) and ZnNPs ($F = 114.22$, $p < 0.0001$), with the interaction between NaCl and ZnNPs also being highly significant ($F = 597.77$, $p < 0.0001$). These results indicate that ZnNPs play a protective role in reducing oxidative stress and lipid peroxidation under salt stress conditions (Fig 10).




Fig. 8: Effect of ZnNPs and salt stress on POD of wheat




Fig. 9: Effect of ZnNPs and salt stress on catalase of wheat




Fig. 10: Effect of ZnNPs and salt stress on MDA of wheat




Fig. 11: Effect of ZnNPs and salt stress on H₂O₂ of wheat

Table 1: Statistical analysis of data showing the all parameters of Wheat (*Triticum aestivum*) by foliar applied zinc oxide nanoparticles under salinity stress

Parameter	NaCl (F-value)	NaCl (P-value)	ZnNPs (F-value)	ZnNPs (P-value)	(P- NaCl * ZnNPs (F-value)	(P- NaCl * ZnNPs (P-value)
Sodium Shoot	2.149	0.1341	171.268	< 0.0001	9.323	< 0.0001
Calcium Root	6.393	0.0049	397.060	< 0.0001	42.184	< 0.0001
Calcium Shoot	14.056	< 0.0001	1231.765	< 0.0001	69.886	< 0.0001
SOD Activity	0.621	0.5437	61.590	< 0.0001	4.509	0.0011
POD Activity	8.781	0.0010	262.243	< 0.0001	60.871	< 0.0001
CAT Activity	2.571	0.0932	70.514	< 0.0001	17.846	< 0.0001
MDA Content	120.746	< 0.0001	114.216	< 0.0001	597.769	< 0.0001
H ₂ O ₂ Levels	11.592	0.0002	471.837	< 0.0001	71.453	< 0.0001

Hydrogen Peroxide (H₂O₂) Levels

ZnNPs application significantly reduced H₂O₂ levels under all salinity treatments, particularly at 250 mg/L ZnNPs under 180 mM NaCl. Statistical analysis ($F = 471.84$, $p < 0.0001$) confirmed the effectiveness of ZnNPs in reducing H₂O₂ accumulation, highlighting their role in enhancing the plant's antioxidant defense system and protecting against oxidative stress (Fig 11, table 1).

4 | DISCUSSION

The results from this study provide valuable insights into the impact of zinc nanoparticles (ZnNPs) on wheat (*Triticum aestivum* L.) under varying levels of salinity stress. Specifically, the study focused on the effects of ZnNPs on sodium (Na⁺) accumulation, calcium (Ca²⁺) uptake, and antioxidant enzyme activities, which are critical to plant growth and stress tolerance. The

findings from the statistical analysis and observed trends highlight the promising role of ZnNPs in mitigating salt stress and improving the physiological response of wheat plants.

Sodium Accumulation in Shoots

Sodium toxicity is a major concern under salt stress, as excessive Na⁺ accumulation in plant tissues can disrupt cellular processes and reduce overall plant productivity (Arif *et al.*, 2020). In this study, Na⁺ levels in wheat shoots were significantly reduced with the application of ZnNPs, particularly at concentrations between 100-250 mg/L. The F-value and P-value from the ANOVA indicate a strong statistical significance in the reduction of Na⁺ accumulation with ZnNPs treatment, confirming their role in mitigating salt-induced sodium toxicity. At higher ZnNP concentrations (200 mg/L and 250 mg/L), a slight reduction in Na⁺ levels was observed, suggesting a threshold beyond which ZnNPs no longer confer additional benefits and may even disrupt ion homeostasis. These results are consistent with previous studies showing that ZnNPs help in regulating ion transport and improving membrane permeability, thus limiting sodium translocation from roots to shoots (Singh *et al.*, 2024).

Calcium Uptake in Roots and Shoots

Calcium plays a vital role in maintaining cell wall stability and activating stress-related signaling pathways (Novaković *et al.*, 2018). In this study, the calcium content in both roots and shoots increased significantly with ZnNPs treatment, particularly at 150 mg/L. This enhancement in calcium uptake was observed under both non-saline and saline conditions, indicating that ZnNPs help mitigate the negative effects of salinity on calcium transport and retention. The highest calcium levels were observed in the roots, with a peak at 150 mg/L ZnNPs, suggesting that ZnNPs may stimulate the calcium uptake pathways in the roots. However, at concentrations above 150 mg/L, calcium levels in both roots and shoots began to decline, suggesting that higher concentrations of ZnNPs may have diminishing returns or cause toxicity. This result supports findings from other studies where ZnNPs were shown to improve calcium uptake under stress conditions, possibly by enhancing the activity of calcium transporters (Ayyaz *et al.*, 2024).

Antioxidant Enzyme Activity (SOD, POD, CAT)

Salt stress induces oxidative stress in plants, leading to the accumulation of reactive oxygen species (ROS), which damage cellular components and impair plant growth (Ahmad *et al.*, 2019). Antioxidant

enzymes such as superoxide dismutase (SOD), peroxidase (POD), and catalase (CAT) play a crucial role in protecting plants from oxidative damage by scavenging ROS (Ighodaro, & Akinloye, 2018). The ZnNPs treatment significantly enhanced the activity of these enzymes, particularly at 200 mg/L ZnNPs, under both moderate and high salinity levels. The F-values and P-values from the statistical analysis demonstrate the strong influence of ZnNPs on boosting antioxidant defense mechanisms. The interaction between NaCl and ZnNPs was highly significant, highlighting that ZnNPs effectively activate antioxidant enzymes under salinity stress. These findings are consistent with previous research, where ZnNPs were shown to increase antioxidant enzyme activity, thus improving plant tolerance to oxidative stress induced by salinity (Aazami *et al.*, 2021).

Malondialdehyde (MDA) and Hydrogen Peroxide (H₂O₂) Levels

MDA is a marker of lipid peroxidation and oxidative damage, while H₂O₂ is a reactive oxygen species (ROS) that accumulates under stress (El-Beltagi, & Mohamed, 2013). In this study, MDA and H₂O₂ levels increased with rising salinity but were significantly reduced with ZnNPs application, especially at 250 mg/L ZnNPs. The reduction in these oxidative stress markers indicates that ZnNPs help in reducing oxidative damage and maintaining cellular integrity under salt stress. Statistical analysis revealed a strong interaction between NaCl and ZnNPs, suggesting that ZnNPs are particularly effective in mitigating oxidative damage under high salinity conditions. The results align with previous studies showing that ZnNPs can improve the antioxidant defense system in plants, which helps in reducing ROS accumulation and limiting oxidative damage (Kumar *et al.*, 2020).

Overall Impact of ZnNPs on Wheat Growth

The application of ZnNPs significantly improved the physiological performance of wheat under salt stress. The enhanced potassium and calcium uptake, along with the increased antioxidant enzyme activity and reduced oxidative stress, suggest that ZnNPs play a crucial role in improving plant tolerance to salinity (Faizan *et al.*, 2021). These results highlight the potential of ZnNPs as a sustainable solution for enhancing nutrient uptake, regulating ion homeostasis, and activating antioxidant defenses, thus improving crop productivity in saline soils.

Conclusion

This study demonstrates the effectiveness of zinc nanoparticles in mitigating salt stress in wheat. The

ZnNPs treatment significantly reduced sodium accumulation in shoots, improved calcium uptake in roots and shoots, and enhanced antioxidant enzyme activity. These findings suggest that ZnNPs can play a vital role in improving wheat growth and productivity under saline conditions, offering a promising strategy for sustainable agriculture in salinity-challenged environments.

Funding: This study was not supported by any public, commercial, or non-profit funding agency.

Conflicts of Interest: The authors confirm no conflicts of interest.

Authors' Contribution: All authors contributed equally to this research work

Generative AI Statements: The authors declare that this manuscript has been written without the use of generative artificial intelligence tools.

Publisher's Note: The content of this article reflects solely the views of the authors and does not necessarily represent the perspectives of their affiliated organizations, the publisher, the editors, or the reviewers. No products or claims discussed are authorized or guaranteed by the publisher.

REFERENCES

- Aazami, M. A., Rasouli, F., & Ebrahimzadeh, A. (2021). Oxidative damage, antioxidant mechanism and gene expression in tomato responding to salinity stress under in vitro conditions and application of iron and zinc oxide nanoparticles on callus induction and plant regeneration. *BMC Plant Biology*, 21(1), 597.
- Ahanger, M. A., Tomar, N. S., Tittal, M., Argal, S., & Agarwal, R. (2017). Plant growth under water/salt stress: ROS production; antioxidants and significance of added potassium under such conditions. *Physiology and Molecular Biology of Plants*, 23(4), 731-744.
- Ahmad, R., Hussain, S., Anjum, M.A., Khalid, M.F., Saqib, M., Zakir, I., Hassan, A., Fahad, S. and Ahmad, S., 2019. Oxidative stress and antioxidant defense mechanisms in plants under salt stress. In *Plant abiotic stress tolerance: Agronomic, molecular and biotechnological approaches* (pp. 191-205). Cham: Springer International Publishing.
- Arif, Y., Singh, P., Siddiqui, H., Bajguz, A., & Hayat, S. (2020). Salinity induced physiological and biochemical changes in plants: An omic approach towards salt stress tolerance. *Plant Physiology and Biochemistry*, 156, 64-77.
- Ayyaz, A., Zhou, Y., Batool, I., Hannan, F., Huang, Q., Zhang, K., Shahzad, K., Sun, Y., Farooq, M.A. and Zhou, W., 2024. Calcium nanoparticles and abscisic acid improve drought tolerance, mineral nutrients uptake and inhibitor-mediated photosystem II performance in *Brassica napus*. *Journal of Plant Growth Regulation*, 43(2), pp.516-537.
- Chattha, M. U., Amjad, T., Khan, I., Nawaz, M., Ali, M., Chattha, M. B., ... & Hassan, M. U. (2022). Mulberry based zinc nano-particles mitigate salinity induced toxic effects and improve the grain yield and zinc bio-fortification of wheat by improving antioxidant activities, photosynthetic performance, and accumulation of osmolytes and hormones. *Frontiers in Plant Science*, 13, 920570.
- Cheeseman, J. (2016). Food security in the face of salinity, drought, climate change, and population growth. In *Halophytes for food security in dry lands* (pp. 111-123). Academic Press.
- Edgerton, M. D. (2009). Increasing crop productivity to meet global needs for feed, food, and fuel. *Plant physiology*, 149(1), 7-13.
- El Sabagh, A., Islam, M.S., Skalicky, M., Ali Raza, M., Singh, K., Anwar Hossain, M., Hossain, A., Mahboob, W., Iqbal, M.A., Ratnasekera, D. and Singhal, R.K., 2021. Salinity stress in wheat (*Triticum aestivum* L.) in the changing climate: Adaptation and management strategies. *Frontiers in Agronomy*, 3, p.661932.
- El-Beltagi, H. S., & Mohamed, H. I. (2013). Reactive oxygen species, lipid peroxidation and antioxidative defense mechanism. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca*, 41(1), 44-57.
- Faizan, M., Bhat, J. A., Chen, C., Alyemeni, M. N., Wijaya, L., Ahmad, P., & Yu, F. (2021). Zinc oxide nanoparticles (ZnO-NPs) induce salt tolerance by improving the antioxidant system and photosynthetic machinery in tomato. *Plant Physiology and Biochemistry*, 161, 122-130.
- Hano, C., & Abbasi, B. H. (2021). Plant-based green synthesis of nanoparticles: Production, characterization and applications. *Biomolecules*, 12(1), 31.
- Heuvelink, E., & Dorais, M. (2005). Crop growth and yield. In *Tomatoes* (pp. 85-144). Wallingford UK: Cabi Publishing.
- Ighodaro, O. M., & Akinloye, O. A. (2018). First line defence antioxidants-superoxide dismutase (SOD), catalase (CAT) and glutathione peroxidase (GPX): Their fundamental role in the entire antioxidant defence grid. *Alexandria journal of medicine*, 54(4), 287-293.
- Kumar, H., Bhardwaj, K., Nepovimova, E., Kuča, K., Singh Dhanjal, D., Bhardwaj, S., Bhatia, S.K., Verma, R. and Kumar, D., 2020. Antioxidant functionalized nanoparticles: A combat against oxidative stress. *Nanomaterials*, 10(7), p.1334.
- Mariani, L., & Ferrante, A. (2017). Agronomic management for enhancing plant tolerance to abiotic stresses—drought, salinity, hypoxia, and lodging. *Horticulturae*, 3(4), 52.
- Novaković, L., Guo, T., Bacic, A., Sampathkumar, A., & Johnson, K. L. (2018). Hitting the wall—Sensing and signaling pathways involved in plant cell wall remodeling in response to abiotic stress. *Plants*, 7(4), 89.
- Okon, O. G. (2019). Effect of salinity on physiological processes in plants. In *Microorganisms in saline environments: strategies and functions* (pp. 237-262). Cham: Springer International Publishing.
- Sharma, K., & Sharma, P. K. (2025). Wheat as a nutritional powerhouse: Shaping global food security. In *Triticum-The Pillar of Global Food Security*. IntechOpen.

- Singh, A., Rajput, V.D., Lalotra, S., Agrawal, S., Ghazaryan, K., Singh, J., Minkina, T., Rajput, P., Mandzhieva, S. and Alexiou, A., 2024. Zinc oxide nanoparticles influence on plant tolerance to salinity stress: insights into physiological, biochemical, and molecular responses. *Environmental Geochemistry and Health*, 46(5), p.148.
- Zhao, L., Lu, L., Wang, A., Zhang, H., Huang, M., Wu, H., ... & Ji, R. (2020). Nano-biotechnology in agriculture: use of nanomaterials to promote plant growth and stress tolerance. *Journal of agricultural and food chemistry*, 68(7), 1935-1947.